

Chapter 13 Exercise 13A

Q. 1. $F = k(l - l_0)$

where l = current length

l_0 = original length

k = spring constant

(i) $F = 10(6 - 5)$

$$\Rightarrow F = 10 \text{ N}$$

(ii) $F = 10(10 - 5)$

$$\Rightarrow F = 50 \text{ N}$$

(iii) $F = 10(5.2 - 5)$

$$\Rightarrow F = 2 \text{ N}$$

Q. 2. $l_0 = 2, k = 9$

(i) $F = 9(3 - 2)$

$$\Rightarrow F = 9 \text{ N}$$

(ii) $F = 9(5 - 2)$

$$\Rightarrow F = 27 \text{ N}$$

(iii) $F = 9\left(\frac{10}{3} - 2\right)$

$$\Rightarrow F = 12 \text{ N}$$

$$kl = F + kl_0$$

$$\Rightarrow 9l = 54 + 9(2)$$

$$\Rightarrow l = 8 \text{ m}$$

Q. 3.



$$F_l = 2(10 - 1) = 18 \text{ N}$$

$$F_r = 4(10 - 1) = 36 \text{ N}$$

$$\therefore F = F_r - F_l = 36 - 18 = 18 \text{ N}$$



$$F_l = 2(x - 1) = 2x - 2$$

$$F_r = 4(20 - x - 1) = 76 - 4x$$

$$F_l = F_r$$

$$\Rightarrow 2x - 2 = 76 - 4x$$

$$\Rightarrow x = 13 \text{ m}$$

from left hand wall (LHW)

Q. 4.



$$F_l = 5(x - 1) = 5x - 5$$

$$F_r = 3(19 - x - 2) = 51 - 3x$$

$$F_l = F_r$$

$$\Rightarrow 5x - 5 = 51 - 3x$$

$$\Rightarrow x = 7 \text{ m from LHW}$$

(ii) $F_l = 5x - 5$


$$F_r = 51 - 3x$$

$$F_r - F_l = 16$$

$$\Rightarrow (51 - 3x) - (5x - 5) = 16$$

$$\Rightarrow x = 5 \text{ m from LHW}$$

Q. 5.



$$F_l = 7(x - 2) = 7x - 14$$

$$F_r = 3(35 - x - 3) = 96 - 3x$$

$$F_l = F_r$$

$$\Rightarrow 7x - 14 = 96 - 3x$$

$$\Rightarrow x = 11 \text{ m from LHW}$$

(ii) $F_l = 7x - 14, F_r = 96 - 3x$

If the force is 40 N to the right, then

$$F_r - F_l = 40$$

$$\Rightarrow (96 - 3x) - (7x - 14) = 40$$

$$\Rightarrow 110 - 10x = 40$$

$$\Rightarrow x = 7$$

$$\Rightarrow 7 \text{ metres from LHW}$$

If the force is 40 N to the left, then

$$F_l - F_r = 40$$

$$\Rightarrow (7x - 14) - (96 - 3x) = 40$$

$$\Rightarrow 10x - 110 = 40$$

$$\Rightarrow x = 15 \text{ m from LHW}$$

Q. 6. $F = k(l - l_0)$

$$= 50(2 - 1)$$

$$= 50 \text{ N}$$

This is the centripetal force and must equal $m\omega^2 r$

$$\therefore m\omega^2 r = 50$$


$$\Rightarrow 1(\omega)^2(2) = 50$$

$$\Rightarrow \omega = 5 \text{ rad/s}$$

FUNDAMENTAL APPLIED MATHEMATICS

Q. 7. $F_{\text{up}} = k(l - l_0)$
 $= 49(x - 1)$
 $= 49x - 49 \text{ N}$

$F_{\text{down}} = mg$
 $= 10(9.8)$
 $= 98 \text{ N}$



But $F_{\text{up}} = F_{\text{down}}$ (in equilibrium)
 $\therefore 49x - 49 = 98$

$$\Rightarrow x = 3 \text{ m}$$

Q. 8. $F_{\text{up}} = k(l - l_0)$
 $= 7(x - 2)$
 $= 7x - 14 \text{ N}$

$F_{\text{down}} = mg$
 $= 2(9.8)$
 $= 19.6$

$F_{\text{up}} = F_{\text{down}}$
 $\Rightarrow 7x - 14 = 19.6$
 $\Rightarrow x = 4.8 \text{ m}$



$F_{\text{up}} = k(l - 0)$
 $= 7(4.8 + x - 2)$
 $= 19.6 + 7x$


$F_{\text{down}} = mg = 2(9.8) = 19.6$

Nett force = $F_{\text{up}} - F_{\text{down}}$
 $= 19.6 + 7x - 19.6 = 7x$

Q. 9. (i) $l^2 = 4^2 + 3^2$
 $\Rightarrow l = 5 \text{ m}$

$\therefore \text{Extension} = 5 - 3$
 $= 2 \text{ m}$

(also $\cos A = \frac{3}{5}$, $\sin A = \frac{4}{5}$)




(ii) **Forces** **Resolved**



$$F = k(l - l_0) = k(5 - 3) = 2k$$

1. $F_{\text{up}} = F_{\text{down}} \Rightarrow \frac{4}{5}F = mg$
 $\Rightarrow \frac{4}{5}(2k) = mg \Rightarrow k = \frac{5mg}{8}$


(iii) 2. Centripetal force = $m\omega^2 r$
 $\Rightarrow \frac{3}{5}F = m\omega^2(3)$
 But $F = 2k = 2\left(\frac{5mg}{8}\right) = \frac{5mg}{4}$
 $\therefore \frac{3}{5}\left(\frac{5mg}{4}\right) = m\omega^2(3)$
 $\Rightarrow \omega^2 = \frac{g}{4} \Rightarrow \omega = \sqrt{\frac{g}{4}} \text{ rads/s}$



$$F_1 = k(l - l_0) = 10(l - 2) = (10l - 20) \text{ N}$$

$$F_2 = k(l - l_0) = 7(l - 1) = (7l - 7) \text{ N}$$

Forces **(Resolved)**



1. $F_{\text{left}} = F_{\text{right}} \Rightarrow F_1 \sin A = F_2 \sin A$
 $\Rightarrow F_1 = F_2 \Rightarrow 10l - 20 = 7l - 7$
 $\Rightarrow l = \frac{13}{3}$
 Now $\sin A = \frac{\text{opp}}{\text{hyp}} = \frac{4}{13} = \frac{12}{13}$


(ii) $\therefore \cos A = \frac{5}{13}$

2. $F_1 = 10l - 20 = \frac{130}{3} - 20 = \frac{70}{3} \text{ N}$


$F_2 = F_1 = \frac{70}{3} \text{ N}$

$F_{\text{up}} = F_{\text{down}} \Rightarrow F_1 \cos A + F_2 \cos A = W$
 $\Rightarrow W = \left(\frac{70}{3}\right)\left(\frac{5}{13}\right) + \left(\frac{70}{3}\right)\left(\frac{5}{13}\right) = \frac{700}{39} \text{ N}$

Q. 11. Forces



Resolved



$$\begin{aligned}
 F_{\text{left}} &= F_{\text{right}} \Rightarrow F_1 \sin \theta = F_2 \sin \theta \Rightarrow F_1 = F_2 \\
 \Rightarrow k_1(l - l_1) &= k_2(l - l_2) \\
 \Rightarrow k_1l - k_1l_1 &= k_2l - k_2l_2 \\
 \Rightarrow l(k_1 - k_2) &= k_1l_1 - k_2l_2 \\
 \Rightarrow l &= \frac{k_1l_1 - k_2l_2}{k_1 - k_2} \\
 \sin \theta &= \frac{\text{opp}}{\text{hyp}} = \frac{d}{l} = \frac{d(k_1 - k_2)}{k_1l_1 - k_2l_2} \quad \mathbf{QED}
 \end{aligned}$$

Exercise 13B

Q. 1. (i) $\frac{2\pi}{7} = \frac{2\pi}{\omega} \Rightarrow \omega = 7$

Max. velocity = $\omega A = 7(5) = 35 \text{ m/s}$

(ii) Max. acceleration = $\omega^2 A = (7)^2(5) = 245 \text{ m/s}^2$

(iii) Total distance covered = $4A = 20 \text{ m}$

Time taken = $\frac{2\pi}{7}$

$$\begin{aligned}
 \text{Average speed} &= \frac{\text{Distance}}{\text{Time}} \\
 &= \frac{20}{\frac{2\pi}{7}} \\
 &= \frac{140}{2\pi} \\
 &= \frac{70}{\pi} \text{ m/s}
 \end{aligned}$$

Q. 2. $v^2 = \omega^2(A^2 - x^2)$

$x = \sqrt{7}, v = 9$

$\Rightarrow 81 = \omega^2(A^2 - 7) \dots \text{Equation 1}$

$x = 2, v = 6\sqrt{3}$

$\Rightarrow 108 = \omega^2(A^2 - 4) \dots \text{Equation 2}$

Dividing equation 1 by 2 gives

$$\frac{81}{108} = \frac{\omega^2(A^2 - 7)}{\omega^2(A^2 - 4)} \Rightarrow \frac{A^2 - 7}{A^2 - 4} = \frac{3}{4}$$

$\Rightarrow 3A^2 - 12 = 4A^2 - 28 \Rightarrow A = 4$

Putting this result into equation 1 gives:

$81 = \omega^2(4^2 - 7) \Rightarrow \omega^2 = 9 \Rightarrow \omega = 3$

Periodic time $T = \frac{2\pi}{\omega} = \frac{2\pi}{3} \text{ s}$

Q. 3. Max. velocity = $\omega A = 6 \dots \text{Equation 1}$

Max. acceleration = $\omega^2 A$

$= 12 \dots \text{Equation 2}$

Dividing equation 2 by equation 1 gives $\omega = 2$

Therefore, $A = 3$

Periodic time $T = \frac{2\pi}{\omega} = \frac{2\pi}{2} = \pi \text{ s}$

To find a when $v = 2\sqrt{5}$:

Step 1: Find x when $v = 2\sqrt{5}$:

$$v^2 = \omega^2(A^2 - x^2)$$

$$\Rightarrow 20 = 2^2(3^2 - x^2)$$

$$\Rightarrow x = \pm 2$$

Step 2: Find a when $x = \pm 2$

$$a = -\omega^2 x = -(2)^2(\pm 2) = \pm 8 \text{ m/s}^2$$

The magnitude of the acceleration is 8 m/s^2

Q. 4. (i) $v^2 = \omega^2(A^2 - x^2)$

$v = 8 \text{ when } x = 1$

$$\Rightarrow 64 = \omega^2(A^2 - 1) \dots \text{Equation 1}$$

$v = 4 \text{ when } x = 7$

$$\Rightarrow 16 = \omega^2(A^2 - 49) \dots \text{Equation 2}$$

Dividing 1 by 2 gives

$$\frac{64}{16} = \frac{\omega^2(A^2 - 1)}{\omega^2(A^2 - 49)}$$

$$\Rightarrow \frac{A^2 - 1}{A^2 - 49} = \frac{4}{1}$$

$$\Rightarrow 4A^2 - 196 = A^2 - 1$$

$$\Rightarrow A = \sqrt{65}$$

(ii) Putting this result into equation 1 gives

$$64 = \omega^2(65 - 1)$$

$$\Rightarrow \omega = 1$$

$$\therefore T = \frac{2\pi}{\omega} = \frac{2\pi}{1} = 2\pi \text{ s}$$

(iii) When $x = 0, v^2 = \omega^2(A^2 - x^2)$

$$= 1(65 - 0) = 65$$

$$\therefore v = \sqrt{65} \text{ m/s}$$

Q. 5. When $x = 1, v = 3, a = 3$

So $v = \omega \sqrt{A^2 - x^2} \dots \textcircled{1}$

and $a = \omega^2 x \dots \textcircled{2}$

$$\Rightarrow 3 = \omega^2(1)$$

$$\Rightarrow \omega = \sqrt{3}$$

\therefore From $\textcircled{1}, 3 = \sqrt{3} \sqrt{A^2 - 1}$

$$\Rightarrow A = 2$$

From $\textcircled{2}, a_{\text{MAX}} = \omega^2 A$

$$\Rightarrow a_{\text{MAX}} = 3(2)$$

$$\Rightarrow a_{\text{MAX}} = 6 \text{ m/s}^2$$

Q. 6. $v^2 = \omega^2(A^2 - x^2)$. But $v = 24$ when $x = 5$.

$$\therefore 576 = \omega^2(A^2 - 25) \dots \text{Equation 1}$$

Also $a = -\omega^2x$. But $a = -20$ when $x = 5$.
(a and x are always of opposite sign)

$$\therefore -20 = -\omega^2(5) \Rightarrow \omega = 2$$

Putting this into equation 1 gives:

$$576 = 4(A^2 - 25) \Rightarrow A^2 - 25 = 144$$

$$\Rightarrow A = 13$$

$$(i) \text{ Amplitude} = A = 13.$$

$$(ii) \text{ Periodic time} = \frac{2\pi}{\omega} = \frac{2\pi}{2} = \pi \text{ s}$$

(iii) In π seconds it performs 1 oscillation.

In 1 second it performs $\frac{1}{\pi}$ oscillations.

In 60 seconds it performs $\frac{60}{\pi} = 19.1$ oscillations.

Answer: 19 complete oscillations.

Q. 7. When $x = \sqrt{2}$, $a = -4\sqrt{2}$, $\therefore a = -\omega^2x$

$$\Rightarrow -4\sqrt{2} = -\omega^2\sqrt{2} \Rightarrow \omega = 2.$$

When $x = \sqrt{2}$, $v = 2$, and $\omega = 2$,

$$\therefore v^2 = \omega^2(A^2 - x^2) \Rightarrow 4 = 4(A^2 - 2)$$

$$\Rightarrow A = \sqrt{3}$$

Start the clock in the centre $\Rightarrow x = A \sin \omega t$
i.e. $x = \sqrt{3} \sin 2t$

To find t when $x = 1.5$: $1.5 = \sqrt{3} \sin 2t$

$$\Rightarrow 3 = 2\sqrt{3} \sin 2t$$

$$\Rightarrow \sin 2t = \frac{3}{2\sqrt{3}} = \frac{\sqrt{3}}{2}$$

$$\Rightarrow 2t = \frac{\pi}{3} \Rightarrow t = \frac{\pi}{6} \text{ s}$$

$F = ma$. But $a = -\omega^2x = -4(1.5) = -6$
and $m = 2 \text{ kg}$.

$$\therefore F = (2)(-6) = -12 \text{ N}$$

The force is of magnitude 12 N.

Q. 8. Amplitude = $\frac{\text{Highest} - \text{Lowest}}{2}$

$$(i) \Rightarrow A = \frac{13 - 3}{2}$$

$$\Rightarrow A = 5 \text{ m}$$

$$(ii) \frac{T}{2} = \frac{11}{2}$$


$$\Rightarrow T = 11 \text{ hrs}$$

$$(iii) T = \frac{2\pi}{\omega}$$

$$\Rightarrow \omega = \frac{2\pi}{T}$$

$$\Rightarrow \omega = \frac{2\pi}{11} \text{ rad/s}$$

Graph:




$$\text{We have } x = -5 \cos \frac{2\pi}{11}t$$

$$\Rightarrow 2 = -5 \cos \frac{2\pi}{11}t$$

$$\Rightarrow t = 3.47 \text{ hrs OR } 3:29 \text{ PM}$$

Q. 9. (i) Consider the Sine Curve:



$$x = A \sin \omega t$$

$$\text{From the graph } \frac{T}{2} = 6 \text{ s}$$

$$\Rightarrow T = 12 \text{ s}$$

$$(ii) \text{ But } T = \frac{2\pi}{\omega}$$

$$\Rightarrow \omega = \frac{2\pi}{12}$$

$$\Rightarrow \omega = \frac{\pi}{6}$$

$$0.6 = A \sin \frac{\pi}{6}(1.5)$$

$$\Rightarrow \frac{3}{5} = A \sin \frac{\pi}{4}$$

$$\Rightarrow \frac{3}{5} = A \frac{1}{\sqrt{2}}$$

$$\Rightarrow A = \frac{3\sqrt{2}}{5} \text{ m}$$

Q. 10. Periodic Time:

$$\frac{T}{2} = 6:58 - 0:58$$

$$\Rightarrow \frac{T}{2} = 6 \text{ hrs}$$

$$\Rightarrow T = 12 \text{ hrs}$$

$$\text{But } T = \frac{2\pi}{\omega}$$


$$\Rightarrow \omega = \frac{2\pi}{12}$$

$$\Rightarrow \omega = \frac{\pi}{6} \text{ rad/hr}$$

Amplitude:

$$A = \frac{18 - 13}{2} \Rightarrow A = 2.5 \text{ m}$$

∴ Mean level (Equilibrium Position) = 15.5 m



6:58 + t is latest time ship can leave.

$$x = A \cos \omega t$$

$$1 = 2.5 \cos \frac{\pi t}{6}$$

$$\Rightarrow t = 2.214 \text{ hrs}$$

$$\Rightarrow t = 2 \text{ hrs } 12' 50.62''$$

$$\text{so } 6:58 + 2:12' 50.62''$$

$$= 9:10' 50.62'' \quad \text{OR} \quad 9:10 \text{ PM}$$

Exercise 13C

Q. 1. $x = 3 \sin 5t$. ∴ $\frac{dx}{dt} = 15 \cos 5t$.

$$\therefore \frac{d^2x}{dt^2} = -75 \sin 5t$$

$$= -25(3 \sin 5t) = -25x$$

Since the acceleration is proportional to x but in the opposite direction, it will perform SHM.

$$x = 3 \sin 5t \Rightarrow 1.5 = 3 \sin 5t$$

$$\Rightarrow \sin 5t = \frac{1}{2} \Rightarrow 5t, \frac{\pi}{6} \Rightarrow t = \frac{\pi}{30} \text{ s}$$

Q. 2. (i) $x = 4 \cos 2t$. ∴ $\frac{dx}{dt} = -8 \sin 2t$

$$\therefore \frac{d^2x}{dt^2} = -16 \cos 2t = -4(4 \cos 2t)$$

$$= -4x$$

Since the acceleration is proportional to the distance from p, but in the opposite direction, it will perform SHM ($A = 4$, $\omega = 2$)

(ii) Greatest distance = $A = 4 \text{ m}$

(iii) Its velocity is zero at the extreme point. Since the clock starts when the particle is at an extreme point, use $x = 4 \cos 2t$.

$$x = 4 \cos 2t \Rightarrow 2.5 = 4 \cos 2t$$

$$\Rightarrow \cos 2t = 0.625$$

$$\Rightarrow 2t = \cos^{-1}(0.625) \Rightarrow 2t = 0.8956$$

$$\Rightarrow t = 0.4478 \text{ s}$$

Q. 3. $x = 9 \cos 3t$

(i) For $t = 0$, $x = 9 \cos 0 \Rightarrow x = 9$

(ii) **Note:** x is measured from equilibrium.

So, when particle has travelled 2 metres, $x = 7$

$$x = 9 \cos 3t$$

$$\Rightarrow 7 = 9 \cos 3t$$

$$\Rightarrow \cos 3t = \frac{7}{9}$$

$$\Rightarrow 3t = \cos^{-1} \frac{7}{9}$$

$$\Rightarrow t = 0.227 \text{ s}$$

Q. 4. (i) $x = 13 \sin(\omega t + \varepsilon)$,

when $t = 0$, $x = 5 \Rightarrow 5 = 13 \sin \varepsilon$

$$\sin \varepsilon = \frac{5}{13} = 0.3846$$

$$\Rightarrow \varepsilon = \sin^{-1}(0.3846) = 0.3948$$

(ii) $v^2 = \omega^2(A^2 - x^2)$, $v = 24$ when $x = 5$.

Also $A = 13$

$$\therefore (24)^2 = \omega^2(13^2 - 5^2)$$

$$\Rightarrow 576 = \omega^2(144) \Rightarrow \omega = 2$$

(iii) $x = 0 \Rightarrow 13 \sin(\omega t + \varepsilon) = 0$

$$\Rightarrow \omega t + \varepsilon = 0 \quad \text{OR} \quad \pi \quad \text{OR} \quad 2\pi \text{ etc.}$$

$$\Rightarrow 2t + 0.3948$$

$$= 0 \quad \text{OR} \quad 3.1416 \quad \text{OR} \quad 6.2832 \text{ etc.}$$

The first time ($t > 0$) will be when

$$2t + 0.3948 = 3.1416$$

$$\Rightarrow 2t = 2.7468$$

$$\Rightarrow t = 1.3734 \text{ s}$$

Q. 5. (i) $x = 3 \cos 2t + 4 \sin 2t$

$$A = \sqrt{3^2 + 4^2} = 5$$

$$T = \frac{2\pi}{2} = \pi$$

(ii) $x = 8 \cos 4t + 6 \sin 4t$

$$A = \sqrt{8^2 + 6^2} = 10$$

$$T = \frac{2\pi}{4} = \frac{\pi}{2}$$

FUNDAMENTAL APPLIED MATHEMATICS

(iii) $x = 12 \cos t + 5 \sin t$

$$A = \sqrt{12^2 + 5^2} = 13$$

$$T = 2\pi$$

(iv) $x = 3 \cos \pi t + \sqrt{7} \sin \pi t$

$$A = \sqrt{3^2 + \sqrt{7}^2} = 4$$

$$T = \frac{2\pi}{\pi} = 2$$

(v) $x = \sin 3t + \cos 3t$

$$A = \sqrt{1^2 + 1^2} = \sqrt{2}$$

$$T = \frac{2\pi}{3}$$

(vi) $x = 21 \sin 2\pi t + 20 \cos 2\pi t$

$$A = \sqrt{21^2 + 20^2} = 29$$

$$T = \frac{2\pi}{2\pi} = 1$$

(vii) $x = \sqrt{3} \sin 5t - \cos 5t$

$$A = \sqrt{\sqrt{3}^2 + 1} = 2$$

$$T = \frac{2\pi}{5}$$

(viii) $x = 2 \sin \frac{t}{2} + 3 \cos \frac{t}{2}$

$$A = \sqrt{2^2 + 3^2} = \sqrt{13}$$

$$T = \frac{2\pi}{\left(\frac{1}{2}\right)} = 4\pi$$

(ix) $x = 24 \sin \frac{t}{4} - 7 \cos \frac{t}{4}$

$$A = \sqrt{24^2 + 7^2} = 25$$

$$T = \frac{2\pi}{\left(\frac{1}{4}\right)} = 8\pi$$

(x) $x = 2 \sin \frac{t}{3} + \cos \frac{t}{3}$

$$A = \sqrt{2^2 + 1^2} = \sqrt{5}$$

$$T = \frac{2\pi}{\left(\frac{1}{3}\right)} = 6\pi$$

Q. 6. $x = 12 \cos t + 35 \sin t$

$$\Rightarrow \dot{x} = -12(1) \sin t + 35(1) \cos t$$

$$\Rightarrow \ddot{x} = -12(1)^2 \cos t - 35(1)^2 \sin t$$

$$\Rightarrow \ddot{x} = -12 \cos t - 35 \sin t$$

(i) $\Rightarrow \ddot{x} = -x$

\Rightarrow SHM with $\omega = 1$ rad/s

(ii) $T = \frac{2\pi}{\omega}$

$$\Rightarrow T = \frac{2\pi}{1}$$

$$\Rightarrow T = 2\pi \text{ s}$$

(iii) $A = \sqrt{12^2 + 35^2}$

$$\Rightarrow A = 37 \text{ m}$$

(iv) For $x = 0$:

$$12 \cos t + 35 \sin t = 0$$

$$\Rightarrow 35 \sin t = -12 \cos t$$

$$\Rightarrow \tan t = \frac{-12}{35}$$

$$\Rightarrow t = -0.3303, -0.3303 + \pi, \text{ etc.}$$

$$\Rightarrow t = -0.3303 + \pi$$

for first positive value.

$$\Rightarrow t = 2.811 \text{ s}$$

Q. 7. $x = 12 \sin 2t + 5 \cos 2t$

$$\dot{x} = 12(2) \cos 2t - 5(2) \sin 2t$$

$$\ddot{x} = -12(2)^2 \sin 2t - 5(2)^2 \cos 2t$$

$$\Rightarrow \ddot{x} = -48 \sin 2t - 20 \cos 2t$$

(i) $\Rightarrow \ddot{x} = -4x$

\Rightarrow SHM with $\omega = 2$ rad/s

(ii) $T = \frac{2\pi}{\omega}$

$$\Rightarrow T = \frac{2\pi}{2}$$

$$\Rightarrow T = \pi \text{ s}$$

(iii) $A = \sqrt{12^2 + 5^2}$

$$\Rightarrow A = 13 \text{ m}$$

(iv) For $x = 0$

$$12 \sin 2t + 5 \cos 2t = 0$$

$$\Rightarrow \sin 2t = \frac{-5}{12} \cos 2t$$

$$\Rightarrow \tan 2t = \frac{-5}{12}$$

$$\Rightarrow 2t = \tan^{-1} \frac{-5}{12} + n\pi \quad \text{for all solutions}$$

$$\Rightarrow 2t = -0.3948 + n\pi$$

$$\Rightarrow t = -0.1974 + \frac{n\pi}{2}$$

$$\Rightarrow n = 1 \text{ gives } t_1 = 1.373 \text{ s}$$

(v) $n = 2 \text{ gives } t_2 = 2.944 \text{ s}$

Q. 8. (i) $x = A \cos (\omega t + \alpha)$

$$\Rightarrow \frac{dx}{dt} = -\omega A \sin (\omega t + \alpha)$$

$$\Rightarrow \frac{d^2x}{dt^2} = -\omega^2 A \cos (\omega t + \alpha) = -\omega^2 x$$

Since the acceleration is proportional to x but in the opposite direction, it will perform SHM.

FUNDAMENTAL APPLIED MATHEMATICS

$$v = -2A \text{ when } x = \frac{3A}{5} \text{ and } A = A$$

$$v^2 = \omega^2(A^2 - x^2)$$

$$\Rightarrow 4A^2 = \omega^2\left(A^2 - \frac{9A^2}{25}\right)$$

$$\Rightarrow 4A^2 = \omega^2\left(\frac{16A^2}{25}\right)$$

$$\Rightarrow \omega^2 = \frac{25}{4}$$

$$\Rightarrow \omega = \frac{5}{2}$$

$$\text{Also } x = \frac{3A}{5} \text{ when } t = 0$$

$$\therefore x = A \cos(\omega t + \alpha)$$

$$\Rightarrow \frac{3A}{5} = A \cos(\alpha)$$

$$\Rightarrow \cos \alpha = \frac{3}{5} = 0.6$$

$$\Rightarrow \alpha = 0.9273 \text{ radians}$$

(ii) Now, $A \cos(\omega t + \alpha) = 0$

$$\Rightarrow \omega t + \alpha = \frac{\pi}{2} \quad \text{OR} \quad \frac{3\pi}{2} \quad \text{OR} \quad \frac{5\pi}{2} \quad \text{etc.}$$

$$\Rightarrow \frac{5}{2}t + 0.9273 = \frac{\pi}{2} = 1.571$$

$$\Rightarrow t = 0.2575 \text{ s}$$

- Q. 9.** Maximum acceleration must be not greater than g if the bodies are to stay on the platform.

$$\Rightarrow \omega^2 A \leq 9.8$$

$$\Rightarrow \omega^2(0.2) \leq 9.8$$

$$\Rightarrow \omega \leq 7.$$

Taking ω at its maximum value, 7.

$$T = \frac{2\pi}{\omega} = \frac{2\pi}{7} = 0.8977 \text{ s}$$

Number of oscillations per minute

$$= \frac{60}{0.8977} = 66 \text{ complete oscillations}$$

- Q. 10.** (a) $x = r \cos \omega t$

$$\therefore \frac{dx}{dt} = -r\omega \sin \omega t$$

$$\therefore \frac{d^2x}{dt^2} = -r\omega^2 \cos \omega t = -\omega^2 x$$

$$\therefore a = -\omega^2 x \therefore \text{SHM}$$



When $x = 0.8$, $v = 6$. When $x = r - 0.2$, $a = -24$ (because a must be negative if x is positive)

$$v^2 = \omega^2(A^2 - x^2).$$

$$\text{Put } x = 0.8, v = 6, A = r$$

$$\Rightarrow 36 = \omega^2(r^2 - 0.64)$$

... Equation 1

$$a = -\omega^2 x.$$

$$\text{Put } x = t - 0.2, a = -24$$

$$\Rightarrow -24 = -\omega^2(r - 0.2)$$

$$\Rightarrow 24 = \omega^2(r - 0.2) \dots \text{Equation 2}$$

Dividing equation 2 by equations 1 gives:

$$\frac{24}{36} = \frac{\omega^2(r - 0.2)}{\omega^2(r^2 - 0.64)}$$

$$\Rightarrow \frac{r - 0.2}{r^2 - 0.64} = \frac{2}{3}$$

(ii) $2r^2 - 1.28 = 3r - 0.6$

$$\Rightarrow 2r^2 - 3r - 0.68 = 0.$$

$$\Rightarrow 200r^2 - 300r - 68 = 0$$

$$\Rightarrow 50r^2 - 75r - 17 = 0$$

$$\Rightarrow (5r + 1)(10r - 17) = 0$$

$$\Rightarrow r = -0.2 \quad \text{OR} \quad r = 1.7$$

$r = -0.2$ has no meaning,
so $r = 1.7$

Putting this result into equation 2
gives: $24 = \omega^2(1.7 - 0.2)$

$$\Rightarrow 24 = \omega^2(1.5) \Rightarrow \omega^2 = 16$$

$$\Rightarrow \omega = 4$$

$$\therefore \text{Period} = \frac{2\pi}{4} = \frac{\pi}{2} \text{ s}$$

- (iii) Start clock at centre: $x = A \sin \omega t$
i.e. $x = 1.7 \sin 4t$

At P_1 , $x = 0.8$

$$\Rightarrow 0.8 = 1.7 \sin 4t$$

$$\Rightarrow \sin 4t = \frac{8}{17} = 0.4706$$

$$\Rightarrow 4t = \sin^{-1}(0.4706) = 0.4900$$

$$\Rightarrow t = 0.1225 \text{ s}$$

At P_2 , $x = 1.5$

$$\Rightarrow 1.5 = 1.7 \sin 4t$$

$$\Rightarrow \sin 4t = \frac{15}{17} = 0.8824$$

$$\Rightarrow 4t = \sin^{-1}(0.8824) = 1.0809$$

$$\Rightarrow t = 0.2702 \text{ s}$$

Time to travel from P_1 to P_2


$$= 0.2702 - 0.1225$$

$$= 0.1477 \text{ s}$$

$$= 0.15 \text{ s}$$

Exercise 13D
Q. 1.

(i)



$$\begin{aligned}F_r &= k(l - l_0) \\&= 2(3 - x - 1) \\&= 4 - 2x\end{aligned}$$

$$\begin{aligned}F_l &= k(l - l_0) \\&= 2(3 + x - 1) \\&= 4 + 2x\end{aligned}$$

$$\begin{aligned}F &= F_r - F_l \\&= 4 - 2x - 4 - 2x \\&= -4x \\F = ma &\Rightarrow -4x = 1(a) \Rightarrow a = -4x\end{aligned}$$

This is SHM with $\omega = 2$.


$$(ii) \text{ Periodic time} = \frac{2\pi}{2} = \pi \text{ s}$$

$$(iii) A = \text{original distance from } 0 = 1.$$

$$\begin{aligned}\text{Midway between the walls} &\Rightarrow x = 0 \\&\Rightarrow v^2 = \omega^2(A^2 - x^2) \Rightarrow v^2 = 4(1 - 0) \\&\Rightarrow v = 2 \text{ m/s}\end{aligned}$$

Q. 2.

(i)



$$\begin{aligned}F_r &= k(l - l_0) \\&= 9(5 - x - 1) \\&= 36 - 9x\end{aligned}$$

$$\begin{aligned}F_l &= 9(5 + x - 1) \\&= 36 + 9x\end{aligned}$$

$$\begin{aligned}F_t &= F_r - F_l \\&= 36 - 9x - 36 - 9x \\&= -18x\end{aligned}$$

$$F = ma$$

$$\begin{aligned}&\Rightarrow -18x = \frac{1}{2}a \\&\Rightarrow a = -36x\end{aligned}$$

It will perform SHM with $\omega = 6$. When it is released, $x = 2$. Therefore $A = 2$.

(ii) It starts from an extreme point.

$$\therefore x = a \cos \omega t \text{ i.e. } x = 2 \cos 6t.$$

We want to find t when $x = 1$.

$$\therefore l = 2 \cos 6t$$

$$\begin{aligned}\Rightarrow \cos 6t &= \frac{1}{2} \\&\Rightarrow 6t = \frac{\pi}{3} \\&\Rightarrow t = \frac{\pi}{18} \text{ s}\end{aligned}$$


$$(iii) v^2 = \omega^2(A^2 - x^2)$$

$$\Rightarrow v^2 = 6^2(2^2 - 1^2) = 108$$

$$\Rightarrow v = \sqrt{108} = 6\sqrt{3} \text{ m/s}$$

Q. 3.

(i)



$$F_r = 20(4 - x - 1)$$

$$= 60 - 20x$$

$$F_l = 20(4 + x - 1) = 60 + 20x$$

$$F = F_r - F_l$$

$$= 60 - 20x - 60 - 20x$$

$$= -40x$$

$$F = ma \Rightarrow -40x = 5a \Rightarrow a = -8x$$

It will perform SHM with $\omega = \sqrt{8}$. When it is released, $x = 3$, therefore $A = 3$.

$$\begin{aligned}(ii) \text{ Maximum speed} &= \omega A = \sqrt{8} (3) \\&= 3\sqrt{8} \text{ m/s.}\end{aligned}$$

$$(iii) \text{ We want to find } x \text{ when } v = \sqrt{8} \text{ m/s.}$$

$$v^2 = \omega^2(A^2 - x^2)$$

$$\Rightarrow 8 = 8(9 - x^2)$$

$$\Rightarrow 9 - x^2 = 1$$

$$\Rightarrow x = \sqrt{8} \text{ m}$$

It starts from an extreme point.

$$\therefore x = A \cos \omega t \Rightarrow x = 3 \cos \sqrt{8}t$$

$$(iv) \text{ To find } t \text{ when } x = \sqrt{8}$$

$$\Rightarrow \sqrt{8} = 3 \cos \sqrt{8}t$$


$$\Rightarrow \cos \sqrt{8}t = \frac{\sqrt{8}}{3} = \frac{2.828}{3} = 0.9427$$

$$\Rightarrow \sqrt{8}t = \cos^{-1}(0.9427) = 0.34$$

$$\Rightarrow t = \frac{0.34}{\sqrt{8}} = 0.12 \text{ s}$$

Q. 4.

(i) Let 0 be the position of equilibrium



$$\begin{aligned}F_r &= k(l - l_0) \\&= 8(7 - d - 1)\end{aligned}$$

$$= 48 - 8d$$


$$\begin{aligned}F_l &= k(l - l_0) \\&= 12(d - 1)\end{aligned}$$

$$= 12d - 12$$

FUNDAMENTAL APPLIED MATHEMATICS

$$\begin{aligned} F_r &= F_l \Rightarrow 48 - 8d \\ &= 12d - 12 \\ \Rightarrow d &= 3 \end{aligned}$$

Answer: 3 metres from left hand wall.



$$\begin{aligned} F_r &= k(l - l_0) \\ &= 8(4 - x - 1) \\ &= 24 - 8x \\ F_l &= k(l - l_0) \\ &= 12(3 + x - 1) \\ &= 24 + 12x \\ F &= F_r - F_l \\ &= 24 - 8x - 24 - 12x \\ &= -20x \\ F &= ma \\ \Rightarrow -20x &= 5a \\ \Rightarrow a &= -4x \end{aligned}$$

It will perform SHM with $\omega = 2$.

- (iii) When it is released its displacement, x , from 0 is $\frac{1}{2}$ m.

Therefore $A = \frac{1}{2}$.

Periodic time $= \frac{2\pi}{\omega} = \frac{2\pi}{2} = \pi$ s

Maximum velocity $= \omega A$

$$\begin{aligned} &= 2\left(\frac{1}{2}\right) \\ &= 1 \text{ m/s} \end{aligned}$$

- (iv) Firstly, find x when $v = \frac{\sqrt{3}}{2}$:

$$v^2 = \omega^2(A^2 - x^2)$$


$$\begin{aligned} \Rightarrow \frac{3}{4} &= 4\left(\frac{1}{2} - x^2\right) \\ \Rightarrow x &= \frac{1}{4} \end{aligned}$$

When $x = \frac{1}{4}$

$$a = \omega^2 x = -(4)\left(\frac{1}{4}\right) = -1 \text{ m/s}^2$$

The acceleration is of magnitude 1 m/s².


$$F = ma \Rightarrow F = (5)(1) = 5 \text{ N}$$



$$\begin{aligned} F_r &= k(l - l_0) \\ &= 9(20 - d - 2) \\ &= 162 - 9d \end{aligned}$$

$$\begin{aligned} F_l &= k(l - l_0) \\ &= 15(d - 2) \\ &= 15d - 30 \end{aligned}$$

$$\begin{aligned} F_r &= F_l \\ \Rightarrow 162 - 9d &= 15d - 30 \\ \Rightarrow d &= 8 \end{aligned}$$



$$\begin{aligned} F_r &= k(l - l_0) \\ &= 9(12 - x - 2) \\ &= 90 - 9x \end{aligned}$$

$$\begin{aligned} F_l &= k(l - l_0) \\ &= 15(8 + x - 2) \\ &= 90 + 15x \end{aligned}$$

$$\begin{aligned} F &= F_r - F_l \\ &= 90 - 9x - 90 - 15x \\ &= -24x \end{aligned}$$

$$\begin{aligned} F &= ma \\ \Rightarrow -24x &= \frac{1}{6}a \\ \Rightarrow a &= -144x \end{aligned}$$

It will perform SHM with $\omega = 12$

- (ii) When it was released its displacement, x , from 0 was 1 m.

Therefore $A = 1$.

$$\begin{aligned} \text{Maximum acceleration} &= \omega^2 A \\ &= 144(1) \\ &= 144 \text{ m/s}^2 \end{aligned}$$

$$\begin{aligned} (\text{iii}) \quad \frac{3}{5} (\text{Max. acceleration}) &= \frac{3}{5}(144) \\ &= \frac{432}{5} \text{ m/s}^2 \end{aligned}$$

When $a = \frac{432}{5}$, what is x ? $a = -\omega^2 x$

$$\begin{aligned} \Rightarrow \frac{432}{5} &= -144x \\ \Rightarrow x &= \frac{-3}{5} \end{aligned}$$

FUNDAMENTAL APPLIED MATHEMATICS

When $x = -\frac{3}{5}$, what is v ?


$$v^2 = \omega^2(A^2 - x^2)$$

$$\Rightarrow v^2 = 144\left(1 - \frac{9}{25}\right)$$

$$\Rightarrow v^2 = 144\left(\frac{16}{25}\right)$$

$$\Rightarrow v = 12\left(\frac{4}{5}\right)$$

$$= \frac{48}{5} = 9.6 \text{ m/s}$$



(i) **Hooke's Law:** $F = k[l - l_0]$

where, k = spring constant

l = current length

l_0 = original length.

At x , NZL: $\Sigma F = ma$

$$\pm F_L = ma$$

$$\Rightarrow -k(l_0 + x - l_0) = ma$$

$$\Rightarrow -5(x) = 20a$$

$$\Rightarrow a = -\frac{x}{4}$$

$$\Rightarrow \text{SHM with } \omega = \frac{1}{2}$$

$$(ii) \Rightarrow T = \frac{2\pi}{\omega} = 4\pi$$

Note: Particle travels from Q to E.P. with SHM. It then travels to P with uniform velocity. (String has gone slack)

$$t_1; \text{ from Q to E.P., is } \frac{T}{4} = \left(\frac{1}{4}\right) \text{ of cycle} \\ = \pi$$


$$t_2; \text{ from E.P. to P, } t = \frac{\text{distance}}{\text{speed}}$$

where $v = \omega A$ at E.P.

$$\Rightarrow t_2 = \frac{l_0}{\left(\frac{1}{2}\right)(1)} = 2$$

so Total Time Taken = $t_1 + t_2 = \pi + 2$,

QED



$$F = k[l - l_0]$$

At position x , NZL: $\Sigma F = ma$

$$\pm -F_L = ma$$

$$\Rightarrow -k(l - l_0) = ma$$

$$\Rightarrow -8[1 - x - 1] = \frac{1}{2}a$$

$$(i) \Rightarrow a = -16x$$

$\Rightarrow \text{SHM, } \omega = 4$

$$\omega = 4, \quad A = 0.5, \quad T = \frac{2\pi}{\omega} = \frac{\pi}{2}$$

$$(ii) \text{ Max } v = \omega A$$

$$\Rightarrow v_{\text{MAX}} = 4\left(\frac{1}{2}\right) \\ = 2 \text{ m/s}$$

$$(iii) x = A \cos \omega t \text{ (Starting at Extreme Point)}$$


Note, when particle has travelled 0.2 m, $x = 0.3$.

$$\Rightarrow 0.3 = 0.5 \cos 4t$$

$$\Rightarrow \cos 4t = \frac{3}{5}$$

$$\Rightarrow 4t = \cos^{-1}\left(\frac{3}{5}\right)$$

$$\Rightarrow t = 0.23 \text{ s}$$



$$(i) \text{ At } x, \text{ NZL: } \Sigma F = ma$$

$$\pm -k(l - l_0) = m\ddot{x}$$

Note: We use \ddot{x} , instead of the more usual a , for acceleration, to avoid confusion. Here A is amplitude, a is original length.

$$\text{So } -k(\ddot{x} + x - a) = k\ddot{x}$$

$$\Rightarrow \ddot{x} = -x$$

$$\Rightarrow \text{SHM with } \omega = 1 \text{ rad/s}$$

$$(ii) t_1, \text{ from } Q \text{ to E.P.} = \frac{T}{4} = \frac{2\pi}{4\omega} = \frac{\pi}{2} \text{ s}$$

$$t_2, \text{ from E.P. to } P, t_2 = \frac{\text{distance}}{\text{speed}}$$


where $V = \omega A$ at E.P.

$$\Rightarrow t_2 = \frac{a}{\omega A} = \frac{a}{1(a)} = 1 \text{ s}$$

$$\therefore \text{Total time} = t_1 + t_2$$

$$= \frac{\pi}{2} + 1 \\ = 2.57 \text{ s}$$

Q. 9.



$$(ii) 1. R = mg = (1)(9.8) = 9.8$$

$$2. F_r = \mu R = \frac{1}{2}(9.8) = 4.9$$

$$3. F_l = k(l - l_0)$$

$$= 7(1.7 + x - 1) \\ = 4.9 + 7x$$

$$F = F_r - F_l$$

$$= 4.9 - 4.9 - 7x$$

$$= -7x$$

$$F = ma$$

$$\Rightarrow -7x = la$$

$$\Rightarrow a = -7x$$

It will perform SHM with $w = \sqrt{7}$.

$$\text{Periodic time} = \frac{2\pi}{\omega} = \frac{2\pi}{\sqrt{7}}$$

(iii) The centre of oscillation, where $x = 0$, is 1.7 m from p . It was released where $x = 2$. Therefore $A = 2$.

It starts from an extreme point

$$\therefore x = A \cos \omega t \Rightarrow x = 2 \cos \sqrt{7}t$$

To find t when $x = 0.3$ (i.e. $2 - 1.7$)

$$0.3 = 2 \cos \sqrt{7}t$$


$$\Rightarrow \cos \sqrt{7}t = 0.15$$

$$\Rightarrow \sqrt{7}t = \cos^{-1}(0.15)$$

$$= 1.4202$$

$$\Rightarrow t = \frac{1.4201}{\sqrt{7}} \\ = 0.54 \text{ s}$$

Q. 10. (i)



$$1. R = mg = 5(9.8) = 49$$

$$2. F_r = R = (1)(49) = 49$$

$$3. F_l = k(l - l_0) \\ = 140(2.35 + x - 2)$$

$$= 49 + 140x$$

$$F = F_r - F_l \\ = 49 - 49 - 140x$$

$$F = ma$$

$$\Rightarrow -140x = 5a$$

$$\Rightarrow a = -28x$$

This is SHM with $\omega = \sqrt{28} = 2\sqrt{7}$.

$$(ii) |QO| = 2.35$$

$$(iii) \text{ It starts when } |qp| = 3$$

$$\therefore 2.35 + x = 3$$

$$\Rightarrow x = 0.65$$

The amplitude is, therefore, 0.65

$$(iv) T = \frac{2\pi}{\omega} = \frac{2\pi}{\sqrt{7}}$$



The journey from A to B can be divided into two parts: A to O and O to B .

$$A \text{ to } O: \frac{1}{4} \text{ of a full cycle } t = \frac{1}{4} \left(\frac{\pi}{\sqrt{7}} \right) \\ = 0.2968$$

O to B : $x = A \sin \omega t$ (starts at centre)

$$0.35 = 0.65 \sin 2\sqrt{7}t$$

$$\Rightarrow \sin 2\sqrt{7}t = \frac{35}{65} = \frac{7}{13} = 0.5385$$

$$\Rightarrow 2\sqrt{7}t = \sin^{-1}(0.5385) = 0.5687$$

$$\Rightarrow t = 0.1075$$


$$\text{Total time} = 0.2968 + 0.1075$$

$$= 0.4043$$

$$= 0.404 \text{ s}$$

Exercise 13E

Q. 1.



(i) At Equilibrium Position (E.P.): $\uparrow = \downarrow$

$$\text{so } k[d - l_0] = 2g$$

$$\Rightarrow 98[d - 1] = 19.6$$

$$\Rightarrow d = 1.2 \text{ m}$$

(ii) NZL: $\Sigma F = ma$

$$\downarrow mg - k[d + x - l_0] = ma$$

$$\Rightarrow 19.6 - 98[0.2 + x] = 2a$$

$$\Rightarrow a = -49x \Rightarrow \text{SHM with } \omega = 7$$

$$(\text{iii}) \Rightarrow T = \frac{2\pi}{\omega} \Rightarrow T = \frac{2\pi}{7} \text{ s}$$

$$(\text{iv}) v_{\text{MAX}} = \omega A \Rightarrow v_{\text{MAX}} = 7(0.2)$$

$$\Rightarrow v_{\text{MAX}} = 1.4 \text{ m}$$


(v) Falls 0.15 metres $\Rightarrow x = 0.2 - 0.15$

$$\Rightarrow x = 0.05$$

We have $x = A \cos \omega t$, (Particle released from extremum position)

$$\text{So } 0.05 = 0.2 \cos 7t \Rightarrow t = 0.188 \text{ s}$$

Q. 2.



At E.P. $\uparrow = \downarrow$

$$\Rightarrow mg = k(d - l_0) \dots (1)$$

$$\Rightarrow 4.9 = 7(d - 1)$$

$$\Rightarrow d = 1.7, A = 1.7 - 1.2 = 0.5$$

NZL: $\Sigma F = ma$

$$\downarrow mg - k[l - l_0] = ma$$

$$mg - k[d + x - l_0] = ma$$

From (1) $k(d - l_0) - k(d - l_0) - kx = ma$

$$(\text{i}) \Rightarrow a = -\frac{7}{0.5}x$$

$$\Rightarrow a = -14x$$

$\Rightarrow \text{SHM}$

$$\Rightarrow \omega = \sqrt{14}$$

$$(\text{ii}) a_{\text{MAX}} = \omega^2 A = 14(0.5)$$

$$\Rightarrow a_{\text{MAX}} = 7 \text{ m/s}^2$$

$$F_{\text{MAX}} = ma_{\text{MAX}} = 0.5(7)$$


$$\Rightarrow F_{\text{MAX}} = 3.5 \text{ N}$$

(iii) When 2 metres below P, $x = -0.3$ with $x = A \cos \omega t$

$$\text{so } -0.3 = 0.5 \cos \sqrt{14}t$$

$$\Rightarrow t = 0.59 \text{ s}$$

Q. 3.



(i) At E.P., $F_{\text{E.P.}} = mg$

$$\Rightarrow k[l_E - l_0] = mg \dots (1)$$

$$\Rightarrow \frac{49}{d} [l_E - d] = mg$$

$$\Rightarrow l_E = \frac{6d}{5}$$

At x, NZL: $\Sigma F = ma$

$$\downarrow mg - F_x = ma$$

$$\Rightarrow mg - k[l_E + x - l_0] = ma$$

$$\Rightarrow mg - mg - kx = ma \quad \text{From (1)}$$

$$\Rightarrow a = -\frac{k}{m}x$$

$$\Rightarrow a = -\frac{49}{d}x$$

$\Rightarrow \text{SHM}$

$$\omega = \frac{7}{\sqrt{d}}$$

$$(\text{ii}) \text{ String becomes slack: } x = -\frac{d}{5}$$

We have $x = A \cos \omega t$

$$\Rightarrow -\frac{d}{5} = \frac{2d}{5} \cos \frac{7}{\sqrt{d}} t$$


$$\Rightarrow \cos \frac{7t}{\sqrt{d}} = -\frac{1}{2}$$

$$\Rightarrow \frac{7t}{\sqrt{d}} = \frac{2\pi}{3}$$

$$\Rightarrow t = \frac{2\pi\sqrt{d}}{21}$$

Note: Hooke's Law: $F = kx$, $x = \text{extension}$.

Q. 4. For the 4 kg mass




For equilibrium $\uparrow = \downarrow$

$$\Rightarrow kx_1 = 4g \quad \text{where } x_1 = \text{extension}$$

$$\Rightarrow k = \frac{4g}{0.08} \Rightarrow k = 490 \text{ N/m}$$

For the combined (6 kg) mass




For equilibrium $\uparrow = \downarrow$

$$\Rightarrow 490x_2 = 6g \quad \textcircled{1}$$

$$\Rightarrow x_2 = 0.12$$

Now, the Amplitude, A is $x_2 - x_1 = 0.04 \text{ m}$

Combined mass at displacement x from equilibrium:



NZL: $\Sigma F = ma$

$$\Rightarrow 6g - k(x_2 + x) = 6a$$

But $6g = 490x_2$ from $\textcircled{1}$

$$\text{(i)} \Rightarrow 490x_2 - 490(x_2 + x) = 6a$$

$$\Rightarrow a = -\frac{245}{3}x$$

So the motion is Simple Harmonic with

$$\omega = \sqrt{\frac{245}{3}}$$


$$\text{(ii)} \Rightarrow T = \frac{2\pi}{\omega}$$

$$= 2\pi \sqrt{\frac{3}{245}} \quad \text{OR} \quad 2\pi \sqrt{\frac{3}{25}} \text{ g s}$$

$$\text{(iii)} \quad V_{\text{MAX}} = \omega A = \sqrt{\frac{245}{3}} (0.04)$$

$$= \frac{7\sqrt{15}}{75} \quad \text{OR} \quad \frac{1}{5}\sqrt{\frac{9}{3}} \text{ m/s}$$

Q. 5. (i) Mass $= V_p = h^3(1000 \text{ s}) = 1000 h^3 \text{ s}$



When it is depressed a distance x , the extra buoyancy B' is given by (note the downward is positive)

$B' = -\text{weight of displaced liquid}$

$$= -V_{pg} = (h^2x)(1000)g = -1000 h^2xg$$

$$F = ma$$

$$\Rightarrow -1000 h^2xg = (1000 h^3)s a$$

$$\Rightarrow a = -\frac{x}{hs}$$

\therefore It will perform SHM with $\omega = \sqrt{\frac{g}{hs}}$.

$$\text{The periodic time} = \frac{2\pi}{\omega} = 2\pi \sqrt{\frac{hs}{g}}$$

$$\begin{aligned} \text{(ii)} \quad \text{In this case } B' &= -(h^2x)(1000 \text{ kg}) \\ &= 1000 h^2xkg \end{aligned}$$

$$F = ma$$

$$\Rightarrow -1000 h^2xkg = 1000 h^3sa$$

$$\Rightarrow a = -\frac{gk}{hs}x$$

\therefore SHM with $\omega = \sqrt{\frac{gk}{hs}}$

$$\text{Periodic time} = \frac{2\pi}{\omega} = 2\pi \sqrt{\frac{hs}{gk}}$$

$$\text{Q. 6.} \quad \text{(i)} \quad \text{As before, 0.6 of its height will be submerged, i.e. } (0.6)(80) = 48 \text{ cm.}$$

$$\text{(ii)} \quad \text{Originally, its displacement from equilibrium is 2 cm} = 0.02 \text{ m.}$$

$$\text{Therefore } A = 0.02 = \frac{1}{50}.$$

$$\text{Mass} = V_p = [(0.8)(0.5)(0.2)] (600) = 48 \text{ kg}$$

When it is displaced a distance x m below the water, the extra buoyancy B' is given by:

$B' = \text{weight of liquid displaced}$

$$= V_{pg} = (0.5)(0.2)(x)(1000)(9.8) = -980x$$

$$F = ma$$

$$\Rightarrow -980x = 48a$$

$$\Rightarrow a = -\frac{980}{48}x = -\frac{245}{12}x$$

This is SHM with $\omega = \sqrt{\frac{245}{12}}$

$$\text{Maximum acceleration} = \omega^2 A$$

$$= \frac{245}{12} \times \frac{1}{50} = \frac{49}{120} \text{ m/s}^2$$

Exercise 13F

$$\begin{aligned} \text{Q. 1. } T &= 2\pi \sqrt{\frac{l}{g}} \\ &= 2\pi \sqrt{\frac{0.5}{9.8}} \\ &= 1.419 \text{ s} \end{aligned}$$

$$\begin{aligned} \text{Q. 2. } T &= 2\pi \sqrt{\frac{l}{g}} \\ \Rightarrow \frac{l}{g} &= \frac{T^2}{4\pi^2} \\ \Rightarrow l &= \frac{gT^2}{4\pi^2} \\ &= \frac{9.8(3)^2}{4\pi^2} \\ \Rightarrow l &= 2.234 \text{ m} \end{aligned}$$

$$\begin{aligned} \text{Q. 3. } T &= 2\pi \sqrt{\frac{l}{g}} \\ &= 2\pi \sqrt{\frac{4}{9.8}} \\ \Rightarrow T &= 4.014 \text{ s} \end{aligned}$$

$$\begin{aligned} \text{Q. 4. } T &= 2\pi \sqrt{\frac{l}{g}} \Rightarrow l = \frac{gT^2}{4\pi^2} \\ \Rightarrow l &= \frac{9.8(5.5)^2}{4\pi^2} \Rightarrow l = 7.51 \text{ m} = 751 \text{ cm} \end{aligned}$$

$$\begin{aligned} \text{Q. 5. } T_1 : T_2 &= 2 : 1 \\ \Rightarrow 2\pi \sqrt{\frac{l_1}{g}} : 2\pi \sqrt{\frac{l_2}{g}} &= 2 : 1 \\ \Rightarrow \sqrt{l_1} : \sqrt{l_2} &= 2 : 1 \\ \Rightarrow l_1 : l_2 &= 4 : 1 \end{aligned}$$

$$\begin{aligned} \text{Q. 6. } l_1 : l_2 &= 4 : 9 \\ \Rightarrow \sqrt{l_1} : \sqrt{l_2} &= 2 : 3 \\ \Rightarrow 2\pi \sqrt{\frac{l_1}{g}} : 2\pi \sqrt{\frac{l_2}{g}} &= 2 : 3 \\ \Rightarrow T_1 : T_2 &= 2 : 3 \end{aligned}$$

Q. 7. Let g' be the acceleration due to gravity at the satellite.
 $g' < g$ (by Newton's Law) $\Rightarrow \frac{l}{g'} > \frac{l}{g}$

$$\Rightarrow 2\pi \sqrt{\frac{l}{g'}} > 2\pi \sqrt{\frac{l}{g}}$$

\Rightarrow its period of oscillation is longer than normal

\therefore it will go slow.

Q. 8. (i) Number of oscillations
 $= 24 \times 60 \times 60 = 86,400$

(ii) The old periodic time = 1

$$\therefore 2\pi \sqrt{\frac{l}{g}} = 1$$

The new periodic time.

$$\begin{aligned} T &= 2\pi \sqrt{\frac{l(1.02)}{g}} \\ &= \sqrt{1.02} \left(2\pi \sqrt{\frac{l}{g}} \right) \\ &= \sqrt{1.02(1)} \\ &= 1.01 \text{ s} \end{aligned}$$

It will now perform $\frac{86,400}{1.01}$ oscillations in a day, i.e. 85,545 oscillations in a day

It performs $(86,400 - 85,545) = 855$ fewer.

Q. 9. Let T = the original time = $\frac{60}{30} = 2 \text{ s}$

Let T' = the new time = $\frac{60}{31} \text{ s}$

$$\begin{aligned} T' : T &= \frac{60}{31} : 2 \\ &= 60 : 62 \\ &= 30 : 31 \end{aligned}$$

$$\begin{aligned} \therefore 2\pi \sqrt{\frac{l'}{g}} : 2\pi \sqrt{\frac{l}{g}} &= 30 : 31 \\ &= l' : l = 30 : 31 \end{aligned}$$

$$\Rightarrow l' : l = 900 : 961$$

$$\Rightarrow \frac{l'}{l} = \frac{900}{961}$$

$$\Rightarrow l' = \frac{900}{961} l = 0.9365l$$

\therefore the % reduction

$$= 100 - 93.65 = 6.35\%$$

Q. 10. Let l' , T' be the new length and periodic time respectively.

$$\begin{aligned} l' : l &= 2 : 1 \therefore T' : T = 2\pi \sqrt{\frac{l'}{g}} : 2\pi \sqrt{\frac{l}{g}} \\ &= \sqrt{2} : \sqrt{1} = 1.414 : 1 \\ \therefore T' &= 1.414 T \end{aligned}$$

There has been an increase of 41% in its periodic time.

FUNDAMENTAL APPLIED MATHEMATICS

- Q. 11.** Let g' = acceleration due to gravity on the moon. l' , T' will be the length and periodic time of the pendulum on the moon.

$$T : T' = 2\pi\sqrt{\frac{l}{g}} : 2\pi\sqrt{\frac{l'}{g'}}$$

$$= 2\pi\sqrt{\frac{8}{l}} : 2\pi\sqrt{\frac{3}{\frac{1}{6}}}$$

$$= \sqrt{8} : \sqrt{18}$$

$$= \sqrt{4} : \sqrt{9}$$

$$= 2 : 3$$

- Q. 12.** (i) As in text.

$$(ii) T = 2\pi\sqrt{\frac{k}{g}}$$

$$\Rightarrow T^2 = 4\pi^2 \frac{k}{g}$$

$$\Rightarrow g = \frac{4\pi^2 k}{T^2} \quad \mathbf{QED}$$

$$\begin{aligned} (iii) \quad & 39 \text{ cycles / min} \\ & = 1 \text{ cycle / } \frac{1}{39} \text{ min} \\ & = 1 \text{ cycle / } \frac{60}{39} \text{ s} \end{aligned}$$

$$\begin{aligned} \text{So, } g &= \frac{4\pi^2(0.6)}{\left(\frac{60}{39}\right)^2} \\ \Rightarrow g &= 10.0 \text{ m/s}^2 \end{aligned}$$

$$\begin{aligned} (iv) \quad \% \text{ Error} &= \frac{(10.0 - 9.8)(100)}{9.8} = 2.04\% \\ &= 2\% \text{ (to nearest percent)} \end{aligned}$$